Review Article

Microbial Biofilms in Persistent Endodontic Infections: Challenges, Novel Disinfection Strategies, and Future Perspectives

Akansha Kashyap[1]
[1]*Dept. of conservative dentistry and endodontics, SGT University, Gurugram.*

Corresponding Author:
Akansha Kashyap
E-mail ID: work.akansha112001@gmail.com

Submission: 03.07.2026 Acceptance: 13.08.2026 Publication: 03.11.2026
https://www.doi.org/10.63778/CJID-DRJPL/2026_03113

Abstract

Persistent endodontic infections represent one of the most clinically challenging scenarios in modern dentistry, primarily attributed to the formation of complex microbial biofilms within the root canal system. These biofilms demonstrate remarkable resistance to conventional chemomechanical instrumentation and irrigant-based disinfection protocols, thereby compromising treatment outcomes. This review comprehensively examines the microbiological basis of biofilm-mediated persistence in root canal infections, with particular emphasis on key pathogenic species such as Enterococcus faecalis, Candida albicans, and Treponema denticola. The structural architecture of endodontic biofilms, the extracellular polymeric substance (EPS) matrix, quorum sensing mechanisms, and phenotypic switching are discussed in depth. Contemporary challenges posed by biofilm communities are critically analyzed alongside novel disinfection strategies including photodynamic therapy (PDT), nanoparticle-based delivery systems, ozone therapy, laser-activated irrigation, and herbal antimicrobials. Future research directions focusing on biofilm-targeted therapeutics, microbiome modulation, and immunological adjuncts are outlined.

Keywords: Endodontic biofilm; Persistent infection; Enterococcus faecalis; Photodynamic therapy; Nanoparticles; Root canal disinfection; Quorum sensing; Extracellular polymeric substance.

Introduction

Endodontic treatment aims to eliminate microorganisms from the infected root canal system and prevent reinfection, enabling periapical tissue healing. Despite advances in endodontic instruments, irrigants, and obturation materials, treatment failure persists in approximately 15-25% of cases, predominantly due to the tenacious nature of microbial biofilms that resist conventional disinfection strategies[(1)].

Biofilms are structured communities of microorganisms enclosed within a self-produced extracellular polymeric substance (EPS) matrix, adhering to surfaces within the root canal system. Within these communities, microbial cells exhibit dramatically altered gene expression, metabolic activity, and phenotypic characteristics compared to their planktonic counterparts. This physiological transformation confers profound resistance to host immune defenses and antimicrobial agents, often 100 to 1,000 times greater than free-floating bacteria[(2)].

The complex anatomy of the root canal system, encompassing lateral canals, isthmuses, apical deltas, and dentinal tubules, creates ecological niches that shelter microbial communities from mechanical and chemical debridement[(3)]. Understanding the biology, ecology, and resistance mechanisms of endodontic biofilms is therefore essential for developing effective therapeutic strategies and improving clinical outcomes.

Microbiology of Endodontic Biofilms

Polymicrobial Nature of Endodontic Infections

Primary endodontic infections are polymicrobial in nature, typically involving 10-30 bacterial species per canal. The microbial community is dominated by obligate anaerobes such as Fusobacterium nucleatum, Prevotella intermedia, Porphyromonas endodontalis, Peptostreptococcus micros, and Treponema denticola. These organisms interact synergistically, forming metabolic networks where the by-products of one species serve as substrates for another, a phenomenon known as syntrophism[(4)].

Molecular techniques including 16S rRNA gene sequencing and next-generation sequencing (NGS) have dramatically expanded our understanding of endodontic microbial diversity, revealing previously unculturable taxa belonging to the phyla Firmicutes, Bacteroidetes, Fusobacteria, Proteobacteria, Spirochaetes, and Synergistetes[(4)].

Key Pathogens in Persistent Infections

Enterococcus faecalis

Enterococcus faecalis is the most clinically significant organism in post-treatment endodontic disease, isolated in 24-77% of failed root canal cases. Its remarkable persistence is attributed to the ability to invade dentinal tubules up to 1,000 μm, survive in alkaline environments (pH 11), endure nutrient starvation, and adhere to dentin collagen via surface

adhesins (Ace, Esp). Its lytic enzymes, including gelatinase, hyaluronidase, and cytolysin, contribute to tissue destruction and immune evasion[5].

Candida Albicans

Candida albicans, a dimorphic fungus, is detected in approximately 7-18% of failed endodontic cases. Its pathogenicity is linked to the yeast-to-hypha transition, penetration of dentinal tubules, co-aggregation with bacteria, and formation of robust biofilms with a thick polysaccharide matrix. The hyphal form enables deeper tissue penetration and greater antifungal resistance, with clinical significance heightened in immunocompromised patients[6].

Treponema denticola and Porphyromonas gingivalis

These obligate anaerobes are critical components of the 'red complex,' frequently associated with periapical abscesses. T. denticola produces proteolytic enzymes (dentilisin, CTLP) and surface lipoproteins that modulate host immune responses, while P. gingivalis secretes gingipains with profound immunomodulatory capacity. Both demonstrate strong synergism within polymicrobial biofilms, enhancing overall virulence of the community[4].

Biofilm Architecture and Development

Biofilm formation is a sequential, highly regulated process occurring in five stages: (1) initial reversible adhesion of planktonic cells, (2) irreversible attachment mediated by adhesin-receptor interactions, (3) microcolony formation through cell proliferation and EPS production, (4) biofilm maturation into a three-dimensional structure with water channels enabling nutrient transport, and (5) dispersal of sessile cells to colonize new sites[2].

The mature biofilm exhibits a characteristic mushroom-like architecture with heterogeneous cell populations embedded in an EPS matrix composed of polysaccharides, proteins, extracellular DNA (eDNA), and lipids. Water channels traversing the matrix facilitate distribution of nutrients and removal of metabolic waste, sustaining deep biofilm layers.

Resistance Mechanisms of Endodontic Biofilms

Extracellular Polymeric Substance (EPS) Matrix

The EPS matrix constitutes 75-90% of biofilm biomass and acts as the primary barrier to antimicrobial penetration. It impedes diffusion of irrigants such as sodium hypochlorite (NaOCl) and chlorhexidine (CHX), binds and inactivates cationic antimicrobials through electrostatic interactions, deactivates reactive oxygen species via enzymatic components (catalase, superoxide dismutase), and provides structural integrity to the community[7].

Quorum Sensing

Quorum sensing (QS) is a cell density-dependent communication system through which bacteria coordinate gene expression via chemical signal molecules called autoinducers (AIs). In endodontic biofilms, QS regulates biofilm initiation and maturation, EPS production, virulence factor secretion, antibiotic resistance gene expression, and biofilm dispersal. Disruption of QS, termed quorum quenching, represents a promising therapeutic target[8].

Persister Cells

Persister cells are a small subpopulation (0.001-1%) of metabolically dormant, phenotypically tolerant bacteria that survive antibiotic exposure without harboring genetic resistance mutations. Upon cessation of antimicrobial treatment, these cells resume normal growth and repopulate the biofilm, contributing to treatment recalcitrance and relapse. E. faecalis is particularly notable for its high persister frequency within endodontic biofilms[5].

Horizontal Gene Transfer

Biofilms provide optimal conditions for horizontal gene transfer (HGT) through transformation, transduction, and conjugation. The close physical proximity of cells and availability of eDNA in the EPS matrix promote exchange of mobile genetic elements including plasmids, transposons, and integrons carrying antibiotic resistance genes, facilitating rapid dissemination of resistance traits[2].

Clinical Challenges in Biofilm Elimination

Root Canal Anatomical Complexity

The root canal system is an intricate anatomical labyrinth comprising lateral canals, apical ramifications, isthmuses, fins, and accessory foramina. Studies employing micro-CT imaging have demonstrated that conventional nickel-titanium rotary instruments leave 35-53% of the root canal surface area untouched during instrumentation, providing protected niches for biofilm communities[3].

Irrigant Limitations

Sodium hypochlorite (NaOCl) at 1-5.25% remains the gold standard endodontic irrigant due to its broad-spectrum antimicrobial activity and tissue-dissolving capacity. However, its efficacy against established biofilms is significantly limited by smear layer obstruction, inactivation by organic matter, inability to penetrate deeply into dentinal tubules, and concentration-dependent cytotoxicity. Chlorhexidine (CHX) at 2% exhibits excellent substantivity but lacks tissue-dissolving ability[9].

Dentinal Tubule Invasion

Bacterial penetration into dentinal tubules is a significant clinical challenge, as organisms such as E. faecalis, Fusobacterium nucleatum, and Actinomyces israelii can invade tubules to depths of 150-1,000 µm. Conventional irrigants, even under agitation, penetrate tubules to a maximum of only 130 µm under clinical conditions, leaving a

substantial zone of bacterial colonization beyond the reach of chemical disinfection[5].

Novel Disinfection Strategies

Photodynamic Therapy (PDT)

Photodynamic therapy (PDT), also termed photoactivated disinfection (PAD), employs a photosensitizer compound, visible light of appropriate wavelength, and molecular oxygen to generate cytotoxic reactive oxygen species (ROS), including singlet oxygen, superoxide radicals, and hydroxyl radicals. These ROS cause irreversible oxidative damage to bacterial cell membranes, proteins, and nucleic acids, resulting in cell death without inducing antimicrobial resistance[10].

Commonly employed photosensitizers in endodontics include toluidine blue O (TBO), methylene blue, chlorophyllin, and curcumin. PDT achieves significant bacterial reduction (2-4 $\log_{10}$ CFU reduction) against E. faecalis biofilms in root canals and can be delivered via thin fiber-optic cables. Limitations include inadequate tissue penetration depth and requirement for specialized equipment.

Nanoparticle-Based Delivery Systems

Nanoparticles offer unique advantages for endodontic disinfection due to their nanoscale dimensions (1-100 nm), high surface-area-to-volume ratio, ability to penetrate dentinal tubules, and capacity for sustained drug release[7]. Key nanoparticle systems include:

- Silver nanoparticles (AgNPs): Broad-spectrum antimicrobial activity through membrane disruption, ROS generation, and inhibition of bacterial enzyme systems. AgNPs at 10-50 µg/mL demonstrate significant reduction of E. faecalis biofilms.
- Zinc oxide nanoparticles (ZnO NPs): Photocatalytic ROS generation with additional anti-biofilm activity independent of light activation, with enhanced efficacy compared to conventional zinc oxide.
- Chitosan nanoparticles: Biopolymer-based cationic nanoparticles with inherent antimicrobial properties capable of encapsulating and delivering antibiotics with controlled release kinetics.
- Calcium hydroxide nanoparticles: Enhanced alkaline antimicrobial activity with improved penetration into dentinal tubules compared to conventional calcium hydroxide pastes.

Ozone Therapy

Ozone (O_3) is a powerful oxidizing agent with broad-spectrum antimicrobial activity achieved through oxidative destruction of bacterial cell membrane lipids, proteins, and DNA. Ozonated water at 4-8 µg/mL has demonstrated equivalent or superior antimicrobial efficacy to 2.5% NaOCl against E. faecalis in root canals in several in vitro studies. Advantages include absence of toxic residues, rapid decomposition into oxygen and water, and non-mutagenicity[8].

Laser-Activated Irrigation (LAI)

Laser-activated irrigation (LAI) enhances the efficacy of conventional irrigants through acoustic streaming and cavitation effects. Photon-Induced Photoacoustic Streaming (PIPS) utilizing sub-ablative Er:YAG laser energy generates powerful acoustic streaming that efficiently distributes irrigant throughout the canal system, including lateral canals and isthmuses, achieving significantly higher bacterial reduction than conventional needle irrigation[10].

Herbal and Plant-Derived Antimicrobials

The emergence of antimicrobial resistance has intensified interest in plant-derived bioactive compounds as alternative or adjunctive endodontic antimicrobials:

- Curcumin: A polyphenol from Curcuma longa with broad-spectrum antimicrobial and anti-biofilm properties; when combined with PDT, achieves superior bactericidal effects against endodontic pathogens.
- Propolis: Contains flavonoids, phenolic acids, and terpenes with significant antimicrobial activity against E. faecalis and Candida albicans.
- Neem (Azadirachta indica): Contains nimbidin and azadirachtin with demonstrated activity against root canal pathogens; 25% neem extract shows efficacy comparable to 2.5% NaOCl in some studies.
- Morinda citrifolia (Noni) juice: Demonstrated efficacy against E. faecalis biofilms in root dentinal tubules.

Passive Ultrasonic Irrigation (PUI)

Passive Ultrasonic Irrigation (PUI) employs ultrasonically activated files or wires to generate acoustic streaming and cavitation in the irrigant within the root canal. The acoustic microstreaming disrupts biofilm architecture, dislodges debris, and enhances irrigant penetration into lateral canals and isthmuses. Studies consistently demonstrate that PUI achieves superior bacterial reduction compared to conventional syringe-needle irrigation[9].

Emerging and Experimental Approaches

Bacteriophage Therapy

Bacteriophages are viruses that specifically infect and lyse bacteria without affecting human cells. Phage therapy offers exquisite bacterial specificity and the capacity to penetrate biofilms through depolymerization of EPS matrices by phage-encoded depolymerases. Phage-antibiotic synergy

(PAS), whereby sub-inhibitory antibiotic concentrations enhance phage infection efficacy, represents a particularly promising therapeutic concept for endodontic biofilm elimination.

Enzyme-Based Biofilm Disruption

Biofilm-dispersing enzymes that degrade EPS matrix components represent targeted anti-biofilm strategies. Deoxyribonuclease I (DNase I) disrupts the eDNA scaffold of the EPS matrix, dispersing established biofilms and enhancing antibiotic penetration. Dispersin B (DspB) degrades poly-N-acetylglucosamine (PNAG), a key EPS component, while proteinase K degrades EPS structural proteins. These enzymes can be incorporated into irrigant solutions or sustained-release intracanal dressings.

Quorum Sensing Inhibitors (QSIs)

Quorum sensing inhibitors represent a paradigm-shifting approach: rather than killing bacteria outright, QSIs disarm them by interfering with intercellular communication, attenuating virulence without imposing selective pressure that drives resistance development. Natural QSIs include furanones (from marine red algae), ajoene from garlic, curcumin, and quercetin. Synthetic QSIs targeting specific autoinducer receptors are under active investigation for endodontic applications.

Future Perspectives

The future management of endodontic biofilm infections is likely to be shaped by several converging scientific and technological developments:

- Personalized endodontics: Metagenomic sequencing may enable individualized antimicrobial regimens tailored to the specific microbial community profile of each patient's infection, moving beyond empirical broad-spectrum treatment.
- Smart nanotechnology: Biofilm-triggered drug release nanoparticles responding to local pH changes, quorum sensing signals, or enzymatic activity will enable precise, on-demand delivery of antimicrobials directly at the infection site.
- Artificial intelligence: Machine learning algorithms integrating CBCT imaging, microbiological data, and clinical parameters may predict treatment outcomes and guide individualized irrigant selection and procedural protocols.
- Anti-biofilm biomaterials: Development of root canal sealers and obturation materials with inherent anti-biofilm surface properties, incorporating antimicrobial peptides or bioactive glass, will reduce post-treatment re-infection risk.
- CRISPR-Cas systems: Programmable DNA-editing technologies offer the potential to specifically target and disrupt virulence genes or antimicrobial resistance determinants within endodontic pathogens without broad-spectrum bactericidal effects.

Conclusion

Microbial biofilms represent the central pathological entity in persistent endodontic infections, conferring extraordinary resistance to conventional chemomechanical disinfection through EPS matrix protection, quorum sensing-mediated community behavior, persister cell formation, and horizontal gene transfer of resistance determinants.

While no single disinfection strategy currently achieves complete biofilm elimination, a multi-modality approach combining thorough chemomechanical preparation with adjunctive techniques such as PDT, laser-activated irrigation, nanoparticle delivery, and ultrasonic agitation represents the most rational clinical strategy. The advancing fields of nanomedicine, synthetic biology, phage therapy, and quorum sensing inhibition offer exciting therapeutic avenues that may fundamentally transform endodontic practice.

A thorough understanding of endodontic biofilm biology is not merely an academic pursuit but an essential foundation for clinical excellence in endodontics. As postgraduate students and future specialists, integration of scientific advances into evidence-based clinical practice is imperative for improving outcomes in patients with persistent endodontic infections.

Source of Support: Nil

Conflict of Interest: Nil

Copyright © 2026 CSMSS Journal of Innovative Dentistry (CJID). This is an open access article, it is free for all to read, download, copy, distribute, adapt and permitted to reuse under Creative Commons Attribution Non Commercial-ShareAlike: CC BY-NC-SA BY 4.0 license.

References

1. Nair PNR. On the causes of persistent apical periodontitis: a review. Int Endod J. 2006;39(4):249-281.
2. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent infections. Science. 1999;284(5418):1318-1322.
3. Vertucci FJ. Root canal morphology and its relationship to endodontic procedures. Endod Topics. 2005;10(1):3-29.
4. Siqueira JF Jr, Rocas IN. Microbiology and treatment of acute apical abscesses. Clin Microbiol Rev. 2013;26(2):255-273.

5. Stuart CH, Schwartz SA, Beeson TJ, Owatz CB. Enterococcus faecalis: its role in root canal treatment failure and current concepts in retreatment. J Endod. 2006;32(2):93-98.

6. Waltimo TM, Orstavik D, Siren EK, Haapasalo MP. In vitro susceptibility of Candida albicans to four disinfectants and their combinations. Int Endod J. 1999;32(6):421-429.

7. Kishen A. Advanced topics in endodontology: nanotechnology and its applications in endodontics. Dent Clin North Am. 2010;54(2):353-364.

8. Nagayoshi M, Kitamura C, Fukuizumi T, Nishihara T, Terashita M. Antimicrobial effect of ozone on bacteria invading dentinal tubules. J Endod. 2004;30(11): 778-781.

9. Zehnder M. Root canal irrigants. J Endod. 2006;32(5):389-398.

10. Bergmans L, Moisiadis P, Van Meerbeek B, Quirynen M, Lambrechts P. Effect of photo-activated disinfection on endodontic pathogens ex vivo. Int Endod J. 2008;41(8):680-689.